



*Anatomy
Passion*



Lecture: CV. S
Done By: Lina Imar




“اللهم لا سهل إلا ما جعلته سهلا وأنت تجعل الصعب إذا شئت سهلا”

General Anatomy

Lecture 14: Cardiovascular System

**Dr. Mohamed Fathi
Elrefai**


**Ass. Professor of Anatomy &
Embryology
mohamed@hu.edu.jo**



THE CARDIOVASCULAR SYSTEM

* **Definition:** It is the major part of the circulatory system.

* It consists of 2 main parts: 1. heart. 2. Blood vessels.



THE HEART

Definition: A conical hollow muscular organ

Function : pumps the blood to various parts of the body.

Site: In the middle mediastinum (middle space of thoracic cavity), between two lungs

Weight: 300 gm in males & 250 gm in females.

Dimensions: 12X9 cm.

Pericardium: membrane that surrounds the heart which is divided into




1) outer fibrous pericardium: thick, fibrous & attached to diaphragm.

2) inner serous pericardium: which consists of 2 layers:

A. outer parietal: lines the fibrous pericardium.


B. inner visceral: covers the heart.

** The space between the 2 layers is called the pericardial cavity.



** **External features of the heart:** it has:


1. Apex.
2. Base.
3. Four borders.
4. Two surfaces
5. Four chambers.



1. The apex of the heart:


Formed of the Lt ventricle & directed downward forward to the left laterally
It lies in the 5th left intercostal space, 9.5 cm from the mid-sternal line.

↪ between the 5th and 6th ribs,





2. The base of the heart:

* Formed of both atria, mainly the left atrium & directed backward, upward & to the right, forming the posterior aspect of the heart.



3. The borders of the heart are:

- Upper border formed by the 2 atria.
- Lower border formed by the 2 ventricles.
- Right border formed by the right atrium.
- Left border formed by the left ventricle & left auricle.



4. The surfaces of the heart:

* It has 2 surfaces:

a. Anterior or sternocostal surface: is divided by coronary sulcus into:

1. Atrial part formed mainly by RT atrium.


2. Ventricular part: subdivided by anterior interventricular sulcus into:

* RT 2/3 formed by RT ventricle. ↪ Main chamber in anterior surface

* LT 1/3 formed by LT ventricle.

↪ A groove that separates the atria from the ventricles

↪ groove that separates the left and right ventricles




b. Inferior or diaphragmatic surface:

* Formed by the 2 ventricles & divided by posterior interventricular sulcus into:

* LT 2/3 formed by LT ventricle. ↪ Main chamber in inferior surface

* RT 1/3 formed by the RT ventricle.



The main chambers in:

The apex: LV
The base: LA

The upper border: LA
The lower border: RV

The right border: RA
The left border: LV

Anterior surface: RV
Inferior surface: LV

5. The chambers of the heart:

a. **The RT atrium:** has 2 walls:

1) Anterior rough or muscular.

2) Posterior smooth receiving the openings of big veins SVC, IVC & Coronary sinus.

b. **The LT atrium:** its wall is mostly smooth, receiving the openings of the 4 pulmonary veins.

c. **The RT ventricle:** is divided into 2 parts:




1) **Rough muscular inflowing part:** receives the blood from the right atrium & contains 3 papillary muscles. → Attached to **Tricuspid**

2) **Smooth outflowing part:** It is called **Infundibulum**: it pushes the blood into the pulmonary trunk.

d. **The LT ventricle:** is divided into 2 parts:


1) **Rough muscular inflowing part:** receives the blood from the left atrium & contains 2 papillary muscles. → Attached to **Bicuspid**

2) **Smooth outflowing part:** It is called **vestibule**: it pushes the blood into the aorta.



The differences between RV and LV

RV	LV
Smooth outflowing = Infundibulum	Smooth outflowing = vestibule
3 papillary muscles	2 papillary muscles
Attached to Tricuspid	Attached to Bicuspid
pushes the blood into the pulmonary trunk.	pushes the blood into the aorta
receives the blood from the right atrium	receives the blood from the left atrium




The valves of the heart:

*There are 2 types of heart valves:

a. Atrio-ventricular (AV) valves: → Separates atrial and ventricles

1. The right valve called Tricuspid. It has 3 cusps separating the right atrium from right ventricle.
2. The left valve called Bicusped or Mitral. It has 2 cusps separating the left atrium from left ventricle.



b. Semilunar valves:

They are formed of 3 cusps, each cusp has a hollow space above called sinus.

1. Aortic valve: has one sinus anterior & 2 sinuses posterior.
2. Pulmonary valve: has one sinus posterior & 2 sinuses anterior.




Blood supply of heart:

oxygenated blood يغذي القلب ب

1. Arterial supply: by the coronary arteries (RT<) that arise from the beginning of ascending aorta.

deoxygenated blood يباخذ من القلب

2. Venous drainage: through small veins that end in short venous channel called coronary sinus that opens in the RT atrium.



Types of Blood Circulations

1. **Systemic circulation**: It is the passage of blood from the Lt ventricle to the whole body via the aorta & its branches then returning the blood back to the Rt atrium through the SVC & IVC.

2. **Pulmonary circulation**: It is the passage of the non-oxygenated blood from the Rt ventricle to the lungs via the pulmonary arteries then returning the blood back to the Lt atrium through the pulmonary veins after oxygenation.

3. **Portal circulation**: It is the passage of the venous blood from the gastrointestinal system carrying the products of digestion & absorption to the liver via the portal vein where the liver cells metabolize these products, then returning back the blood to the systemic circulation through the hepatic veins to the IVC to the right atrium.


Systemic circulation



Pulmonary circulation





Portal circulation



**Sinusoid has the main role in portal circulation



type of small blood vessel found in certain tissues, particularly in the liver it plays important roles in the exchange of nutrients, gases, and waste products between the blood and surrounding tissues.



BLOOD VESSELS

* Types of blood vessels:

1. Arteries. الشرايين
2. veins. الأوردة
3. Capillaries. الشعيرات الدموية





The blood leaves the left ventricle of the heart through the **aorta**, which divides into **smaller branches** to supply the different systems & tissues of the body.

These branches divide into **smaller & smaller arteries** till they end inside tissues by giving small **arterioles**. These arterioles divide into **smaller arterioles** & finally, the **smaller ones join the capillaries**.

Capillaries collect into very minute **venules**, which **collect into small veins**. These veins **unite to form large veins**, which **collect into larger veins**, & finally, **veins collect in 2 big veins**; Superior vena cava (SVC) & inferior vena cava (IVC), which open into the right atrium of the heart.

Heart → Aorta → smaller branches → arterioles → smaller arterioles → capillaries → venules → small veins → larger veins → IVC, SVC → RA

Arteries	Veins
1. Carry blood away from the heart.	1. Carry blood towards the heart.
2. Carry oxygenated bl. (except Pulmonary A.)	2. Carry deoxygenated bl. (except Pulmonary V.)
3. Divide into branches.	3. Collect from tributaries.
4. Its wall is rich in smooth ms. & elastic fibers (Non-compressible).	4. Its wall is poor in its smooth muscle & elastic fibers (compressible).
5. Thick-walled & narrow lumen.	5. Thin-walled & wide lumen.
6. Do not contain valves.	6. Contain valves.



Arterial Anastomosis

* This is a connection between branches of one artery & branches of adjacent artery.

* It allows a collateral circulation in case of obstruction of one of the 2 arteries.

يعني لو في شريان سكر ف هون المفروض انه خلص يبطل يوصل دم لباقي المناطق بس لانه في هاي الوصلات بينهم ف حيصير في شرايين بديلة فالدم حيكمل فيهم لباقي المناطق

alternative routes of blood flow that can develop in response to a blockage or narrowing of a blood vessel.

* It is present mainly around joints.

* Arteries which do not anastomose are called end arteries.




End Arteries

* Are of 2 types:

1. **Anatomical end arteries:** which do not acquire any sort of anastomosis e.g. central retinal artery. يعني لو الشريان سكر فالشخص سيفقد البصر تماما لأنه ما في شريان بديل

2. **Functional end arteries:** which have some anastomosis between its arteries, however, it is insufficient to compensate the obstructed artery, e.g. coronary arteries of the heart. يعني هون في بديل لكنه مش كافي في حال صار في انسداد للشريان الأساسي



Types of anastomosis between arteries & veins

A. Capillaries: Small vascular connections present in all body organs connecting the small arterioles to the small venules.

B. Direct arterio-venous Shunt:

* This is a direct shunt between arteries & veins in the tissues.

* It is present in certain areas, such as in palm of hand, sole of foot, auricle, nose, lips & gastro-intestinal tract.




* It plays an important role in:

1. Regulation of the blood flow to each organ.

2. Regulation of body temperature (since it can help losing temperature in some cases or storing temperature in other cases).

C. Sinusoids: are wide tortuous vascular spaces lined with phagocytes, present in liver, spleen & bone marrow. They slow down blood flow allowing maximum exchange of O₂, CO₂ & nutrients between blood & tissues.

D. Cavernous (erectile) tissue: Small vascular spaces filled with blood, present in the erectile tissues of penis & clitoris.





Tortuous arteries

* Are irregular arteries & are present in:


a. Movable organ: as facial Artery

b. Expansile organ: as arteries of uterus & urinary bladder.

c. Protrudable organ: as lingual Artery



Main Arteries of the Body





* Important arteries originating from heart:

I. Pulmonary Trunk: arises from right ventricle carrying deoxygenated blood to lungs.

II. Aorta: arises from left ventricle carrying oxygenated blood to be distributed to all systems of body.

The AORTA




* It is divided into 4 parts: **ascending aorta**, **aortic arch**, **descending thoracic aorta** & **abdominal aorta**.

1. Ascending Aorta:

* It passes upward to the right within the pericardium.

* It gives the **RT & LT coronaries** (which supply the heart).



2. Aortic Arch:

* It lies within the superior mediastinum of the thoracic cavity in front of trachea & esophagus.

* It gives 3 branches:

1. **Left subclavian Artery** : which lies in neck & continues in left upper limb as left axillary artery.

2. **Left common carotid Artery** : which supplies left half of head & neck.

3. **Brachiocephalic (Innominate) Artery** : which divides into:


A) **Right Subclavian Artery** : which lies in neck & continues in right upper limb as right axillary Artery

B) **Right Common carotid Artery** : which supplies right half of head & neck.

* Each common carotid Artery divides into:

a) **External carotid Artery** : which supplies mainly the structures of head & neck outside the skull.

b) **Internal carotid Artery** : which enters the skull & supplies mainly brain and intracranial structures.



Main arteries of upper limb

1) **Subclavian artery** : Continues in the upper limb as axillary artery.


2) **Axillary artery** : continues as Brachial artery in the arm.

3) **Brachial artery** : descends to the cubital fossa (in front of elbow), where it lies medial to the tendon of biceps muscle. This is an important site, because we put the stethoscope on it when we measure the blood pressure.

* One cm below bent of elbow, the brachial artery divides into:


A) **Ulnar artery** : which runs along the medial side of forearm.

B) **Radial artery** : which runs along the lateral side of forearm. Above wrist, we can feel the pulse of the radial artery lateral to the tendon of flexor carpi radialis.




3. Descending Thoracic Aorta:

- * It lies in the posterior mediastinum.
- * It starts at the level of T4 & ends at the level of T12.
- * It gives 2 groups of branches:
 - Parietal group** → supplying the thoracic cage: e.g.: Intercostal arteries.
 - Visceral group** → supplying the lungs, trachea & esophagus.




4. Abdominal Aorta:

- * Lies in front of lumbar vertebrae.
- * It starts at T12 & ends at L4.
- * It gives 3 groups of branches:
 - Single branches:** from its anterior aspect for gastrointestinal tube & its related glands (liver, pancreas & spleen) ; They are:
 - 1) **celiac trunk.** 2) **Superior mesenteric Artery** 3) **Inferior mesenteric Artery**



- Left & right**
- Paired branches:** from its lateral aspect.


1. **Phrenic artery:** to diaphragm.
2. **Middle suprarenal artery:** to suprarenal gland.
3. **Renal artery:** to kidney.
4. **Gonadal artery:** testicular artery (to testis) or ovarian artery (to ovary).
5. **Four Lumbar arteries:** to abdominal wall.




- Terminal branches:** Two common iliac arteries

Common Iliac Arteries


- * Each common iliac artery divides into:
 1. **Internal iliac artery:** which is responsible for blood supply of pelvic organs.
 2. **External iliac artery:** which continues in thigh & forms femoral artery, which ends in lower one third of thigh by forming popliteal A. which runs in the back of the knee and ends by dividing into:
 - a. **anterior tibial Artery:** in front of leg.
 - b. **posterior tibial Artery:** in back of leg.



The AORTA



Main Veins of the Body



** There are 2 types of veins in the body:

1. The superficial veins: lie beneath the skin.

2. The deep veins: accompany the arteries. Some small arteries are accompanied by two veins called venae comitants.

1. Veins of the heart end in the coronary sinus.



2. Veins from the head & neck are collected into internal jugular vein.

3. Veins of the upper limb are collected in the subclavian vein.

** Internal jugular unites with the subclavian vein to form brachiocephalic vein.

** The 2 brachiocephalic veins unite to form the SVC.


4. Veins of the abdomen, pelvis & lower limbs end in the IVC.



Superficial veins used in intravenous injection or taking blood sample:

1. Cephalic vein: Starts at the lateral end of dorsal venous arch.

2. Basilic vein: Starts at the medial end of the dorsal venous arch.



* The cephalic & basilic veins are joined at the elbow by a vein called median cubital vein. This vein is the commonest vein used for intravenous injection.

Types of CIRCULATION

• **Coronary circulation** - the circulation of blood within the heart.


• **Pulmonary circulation** - the flow of blood between the heart and lungs.

• **Systemic circulation** - the flow of blood between the heart and the cells of the body.


• **Portal circulation** - the flow of blood between two set of capillaries before draining in systemic veins.

• **Fetal Circulation**


CORONARY CIRCULATION: ARTERIAL SUPPLY



SYSTEMIC AND PULMONARY CIRCULATION





PORTAL CIRCULATION



FETAL CIRCULATION

allows the fetus to receive oxygen and nutrients from the mother's blood and dispose of waste products without having to rely on its own lungs



FETAL CIRCULATION

placenta → Umbilical vein → IVC [through Ductus venosus] → RA → LA [Through Foramen Ovale] → RV → pulmonary trunk

مش مطلوبة هاي التفاصيل، المطلوب نعرف انها دورة بين دم الام ودم الجنين وانه المشيمة بتسوي زي الرئة بتحول الدم غير المؤكسد اللي طالع من الجنين لدم مؤكسد بيرجع للجنين مرة تانية

1) the space between the outer parietal layer and the visceral layer of the heart is called

- A) pleural cavity
- B) pericardial cavity
- C) fibrous pericardium
- D) myocardium

2) the apex of the heart is formed and located and directed in :

- A) left ventricle, downward backward to the left, in the 5th intercostal space
- B) right ventricle, downward forward to the left, in the 5th intercostal space
- C) left ventricle, upward forward to the left, in the 5th intercostal space
- D) left ventricle, downward forward to the left, in the 5th intercostal space

3) the main chamber that form the diaphragmatic surface of the heart is :

- A) RT 2/3 formed by RT ventricle
- B) RT 1/3 formed by the RT ventricle
- C) LT 2/3 formed by LT ventricle
- D) LT 1/3 formed by LT ventricle

4) The RT atrium chamber receive all of these opening except

- A) pulmonary veins
- B) superior vena cava
- C) inferior vena cava
- D) coronary sinus

5) the Aortic valve is a Semilunar valves that has one sinus anterior and 2 sinuses posterior

- A) True
- B) False

6)all of these sentences is correct except :

- A)In the RT ventricle chamber, the Rough muscular inflowing part receives the blood from the right atrium and contains 3 papillary muscle
- B)In the LT ventricle chamber, the Rough muscular inflowing part receives the blood from left atrium and contains 2 papillary muscles
- C)In the RT ventricle chamber, the Smooth outflowing part is called vestibule and pushes the blood into the pulmonary trunk
- D)In the LT ventricle chamber, the Smooth outflowing part pushes the blood into the aorta

7)which of these statement is correct related to pulmonary circulation:

- A)It is the passage of blood from the Lt ventricle to the whole body via the aorta
- B)It is the passage of the non-oxygenated blood from the Rt ventricle to the lungs via the pulmonary veins then returning the oxygenated blood to the Lt atrium through puronar arteries
- C)It is the passage of the non-oxygenated blood from the Rt ventricle to the lungs via the pulmonary arteries then returning the oxygenated blood to the Lt atrium through pulmonary veins
- D)non of the above

8)the correct order of the passage of the blood through the body is:

- A)LT atrium - LT ventricle -- aorta -- small branches - arteries -- arterioles - capillaries -- venules -- veins -- IVC or SVC --- RT atrium
- B)LT ventricle -- LT atrium -- aorta -- small branches - arteries - arterioles --capillaries -- venules -- veins - IVC --- SVC -- RT atrium
- C)LT atrium - LT ventricle - pulmonary arteries --- small branches - arteries --arterioles - capillaries - venules -- veins --- IVC -- SVC - RT atrium
- D)RT atrium --RT ventricle -- aorta -- small branches -- arteries - arterioles - capillaries -- venules - veins -- IVC --- SVC --- LT atrium

9)the pulmonary veins carry deoxygenated blood and pulmonary arteries carry the oxygenated

- A)True.
- B)False

10)All of these are true about The Direct arterio-venous Shunt except :

- A)is a direct shunt between arteries &veins in the tissues
- B)present in all body organs
- C)It plays an important role in Regulation of the blood flow
- D)It is present in certain areas, such as in palm of hand

11)all of these are tortuous arteries except:

- A)facial A
- B)arteries of uterus
- C)lingual A
- D)femoral artery

12)the left subclavian artery is :

- A)a branch of Aortic arch that supplies left half of head & neck.
- B)a branch of Aortic arch which lies in neck & continues in left upper limb as left axillary artery.
- C)a branch of abdominal aorta
- D)Continues in the upper limb as Brachial artery

13)all of theses are features of veins except

- A)Carry blood towards the heart
- B)Collect from tributaries.
- C)Thin-walled & narrow lumen
- D)Its wall is compressible

14)The Descending Thoracic Aorta supplied all of the following except

- A)thoracic cage
- B)lungs
- C)intracranial structure
- D)trachea

15)the incorrect pairs regarding the branches of aorta of the following is

- A)Descending Thoracic Aorta -- starts at the level of T4 & ends at the level of T12
- B)testicular artery - to ovary
- C)External carotid A - supplies the structures of head & neck outside the skull
- D)Brachial artery - we put the stethoscope on it when we measure the blood pressure.

16)choose the correct statement regarding the base of the heart :

- A)Formed of both atrium, backwards forward to the right
- B)Formed of both ventricle, backwards upward to the right
- C)Formed of both atrium, backwards upward to the right
- D)Formed of both atrium, backwards upward to the left

17)The vein is the commonest vein used for intravenous injection :

- A)median cubital vein
- B)brachiocephalic veins
- C)internal jugular vein.
- D)subclavian vein

Answers:

- 1)B
- 2)D
- 3)C
- 4)A
- 5)A
- 6)C
- 7)C
- 8)A
- 9)B
- 10)B
- 11)D
- 12)B
- 13)C
- 14)C
- 15)B
- 16)C
- 17)A

Done by: Lina Imar