



تَوِير

BIOLOGY

Lec no :

File Title: Chapter 13

Done By: Leen Abu Salma

وقل صرنا زكياتا علمنا



Overview: Variations on a Theme

- Living organisms are distinguished by their ability to reproduce their own kind
- **Genetics** علم الوراثة is the scientific study of heredity and variation
- **Heredity** الوراثة is the transmission of traits صفة وراثية from one generation to the next
- **Variation** التنوع is demonstrated by the differences in appearance ظهرت that offspring الذرية / النسل show from parents and siblings

Figure 13.1



Concept 13.1: Offspring acquire genes from parents by inheriting chromosomes

- In a literal sense, children do not inherit particular physical traits from their parents
- It is genes that are actually inherited

Inheritance of Genes

- ^{وحدة الموراثية} **Genes** are the units of heredity, and are made up of ^{شرايط} segments of DNA
- Genes are passed to the next generation via reproductive cells called **gametes** (sperm and eggs)
- Each gene has a specific location called a **locus** on a certain chromosome
- Most DNA is packaged into chromosomes

Gene موقع ال
Chromosome على ال


Comparison of Asexual and Sexual Reproduction

- In **asexual reproduction**, a single individual passes genes to its offspring without the fusion of gametes
- A **clone** ^{نسخ} is a group of genetically identical individuals from the same parent
- In **sexual reproduction**, two parents give rise to offspring that have unique combinations of genes inherited from the two parents



Video: Hydra Budding

Figure 13.2




(a) Hydra

© 2011 Pearson Education, Inc.
*asexual reproduction
(mitosis)*



(b) Redwoods

Figure 13.2a



(a) Hydra

© 2011 Pearson Education, Inc.

Figure 13.2b



(b) Redwoods

© 2011 Pearson Education, Inc.

الإخصاب

Concept 13.2: Fertilization and meiosis alternate in sexual life cycles

دورة الحياة

- A **life cycle** is the generation-to-generation sequence of stages in the reproductive history of an organism

Sets of Chromosomes in Human Cells

*(mitosis occurs)
nonreproductive cells*

- Human **somatic cells** (any cell other than a gamete) have 23 pairs of chromosomes
- A **karyotype** ^{البصمة الوراثية} is an ordered display of the pairs of chromosomes from a cell
- The two chromosomes in each pair are called **homologous chromosomes**, or homologs
- Chromosomes in a homologous pair are the same length and shape and carry genes controlling the same inherited characters

Figure 13.3

APPLICATION




* maternal → From the mother

* paternal → From the father

* synapsis: pairing of homologous chromosomes.

TECHNIQUE

Pair of homologous duplicated chromosomes



5 μ m





Figure 13.3a



© 2011 Pearson Education, Inc.

Figure 13.3b

Pair of homologous duplicated chromosomes



5 μm





Figure 13.3c



- The **sex chromosomes**, which determine the sex of the individual, are called **X** and **Y**
- Human females have a homologous pair of X chromosomes (**XX**)
- Human males have one **X** and one **Y** chromosome (**X Y**)
- The remaining 22 pairs of chromosomes are called **autosomes**

– *Autosomes: carry genes for body traits.*

- Each pair of homologous chromosomes includes one chromosome from each parent
- The 46 chromosomes in a human somatic cell are two sets of 23: one from the mother and one from the father
- A **diploid cell** ($2n$) has two sets of chromosomes
- For humans, the diploid number is 46 ($2n = 46$)

$$1n = 23$$

* A Haploid cell ($1n$)

- In a cell in which DNA synthesis has occurred, each chromosome is replicated
- Each replicated chromosome consists of two identical sister chromatids

Figure 13.4


* gonads : هي خلية تتواجد في الأعضاء التناسلية.

Key

- $2n = 6$ {
- **Maternal set of chromosomes ($n = 3$)**
 - **Paternal set of chromosomes ($n = 3$)**

* gonads → male (testis)
→ female (ovaries)

Sister chromatids of one duplicated chromosome



Centromere

Two nonsister chromatids in a homologous pair

Pair of homologous chromosomes (one from each set)

cohesin protein

* بصطفا على شكل رباعي tetrad


- A gamete (sperm or egg) contains a single set of chromosomes, and is **haploid** (n)
- For humans, the haploid number is 23 ($n = 23$)
- Each set of 23 consists of 22 autosomes and a single sex chromosome
- In an unfertilized egg (ovum), the sex chromosome is X
- In a sperm cell, the sex chromosome may be either X or Y

Behavior of Chromosome Sets in the Human Life Cycle

- **Fertilization** is the union of gametes (the sperm and the egg)
- The fertilized egg is called a **zygote** and has one set of chromosomes from each parent
- The zygote produces somatic cells by mitosis and develops into an adult

- At sexual maturity, the ovaries and testes produce haploid gametes
- Gametes are the only types of human cells produced by **meiosis**, rather than mitosis
- Meiosis results in one set of chromosomes in each gamete
- Fertilization and meiosis alternate in sexual life cycles to maintain chromosome number

Figure 13.5



The Variety of Sexual Life Cycles

- The alternation of meiosis and fertilization is common to all organisms that reproduce sexually
- The three main types of sexual life cycles differ in the timing of meiosis and fertilization



- 
- Gametes are the only haploid cells in animals
 - They are produced by meiosis and undergo no further cell division before fertilization
 - Gametes fuse to form a diploid zygote that divides by mitosis to develop into a multicellular organism

Figure 13.6


Key

- Haploid (n)
- Diploid ($2n$)




(a) Animals

© 2011 Pearson Education, Inc.



(b) Plants and some algae




(c) Most fungi and some protists

Figure 13.6a

Key

- Haploid (n)
- Diploid ($2n$)



(a) Animals

- Plants and some algae exhibit an **alternation of generations**
- This life cycle includes both a diploid and haploid multicellular stage
- The diploid organism, called the sporophyte, makes haploid spores by meiosis



- 
- Each spore grows by mitosis into a haploid organism called a gametophyte
 - A gametophyte makes haploid gametes by mitosis
 - Fertilization of gametes results in a diploid sporophyte

Figure 13.6b

Key

- Haploid (n)
- Diploid ($2n$)



(b) Plants and some algae



- 
- In most fungi and some protists, the only diploid stage is the single-celled zygote; there is no multicellular diploid stage
 - The zygote produces haploid cells by meiosis
 - Each haploid cell grows by mitosis into a haploid multicellular organism
 - The haploid adult produces gametes by mitosis

Figure 13.6c

Key

→ Haploid (n)

→ Diploid ($2n$)



(c) Most fungi and some protists

- Depending on the type of life cycle, either haploid or diploid cells can divide by mitosis
- However, only diploid cells can undergo meiosis
- In all three life cycles, the halving and doubling of chromosomes contributes to genetic variation in offspring

in sex organs

Concept 13.3: Meiosis reduces the number of chromosome sets from diploid to haploid

- Like mitosis, meiosis is preceded by the replication of chromosomes
- Meiosis takes place in two sets of cell divisions, called **meiosis I** and **meiosis II**
- The two cell divisions result in four daughter cells, rather than the two daughter cells in mitosis
- Each daughter cell has only half as many chromosomes as the parent cell

The Stages of Meiosis

- After chromosomes duplicate, two divisions follow
 - ① *halving the number of chromosomes* Meiosis I (reductional division): homologs pair up and separate, resulting in two haploid daughter cells with replicated chromosomes
 - ② Meiosis II (equational division) sister chromatids separate (*Very similar to mitosis*) .
- The result is four haploid daughter cells with unreplicated chromosomes

Figure 13.7-1

Interphase





Figure 13.7-2

Interphase



Meiosis I






Figure 13.7-3

Interphase




cell cycle
for replication

Meiosis I (reductional division) *انقسام اختزالي*



Meiosis II (equational division) *انقسام مساوي*



gametes

- Meiosis I is preceded by interphase, when the chromosomes are duplicated to form sister chromatids
- The sister chromatids are genetically identical and joined at the centromere
- The single centrosome replicates, forming two centrosomes



BioFlix: Meiosis

- Division in **meiosis I** occurs in four phases


① Prophase I

② Metaphase I

③ Anaphase I

④ Telophase I and cytokinesis

Figure 13.8



* Prophase 1

- Centrosomes start moving to the opposite pole of the cells and while they are moving they start releasing spindle microtubules.
- nucleolus will disappear.
- nuclear envelope will disappear.
- each chromosome will appear as a two sister chromatids.
- the chromosomes will be thicker and shorter and appear under the microscope.
- ^(pairing) synapsis of homologous chromosomes.
- Interaction between two non-sister chromatids.
- The chiasmata: When the non-sister chromatids cross with each other they will exchange of genetic material, and we call this process "crossing over".

* Metaphase 1

- Centrosomes will settle at the opposite poles and the spindle formation is completed.
- lining for homologous pairs of chromosomes on metaphase plate, each chromosome is attached to kinetochore microtubule.

* Anaphase 1


- inactivation for cohesin protein.
- shortening and contraction of kinetochore microtubules.
- separation of each of the homologous chromosomes, each homologous pair will move to the opposite pole of the cell.
- elongation of the cell.

* Telophase 1

- Reformation of the nuclei.
- cytokinesis occur.

- 2 cells each cell have half number of the chromosomes.
- These 2 cells are non-identical to each other.

Figure 13.8a



Prophase I

- Prophase I typically occupies more than 90% of the time required for meiosis
- Chromosomes begin to condense
- In **synapsis**, homologous chromosomes loosely pair up, aligned gene by gene

- In **crossing over**, nonsister chromatids **exchange** DNA segments
- Each pair of chromosomes forms a tetrad, a group of four chromatids ^{وباعيات}
- Each tetrad usually has one or more **chiasmata**, X-shaped regions where crossing over occurred

Metaphase I

- In metaphase I, tetrads line up at the metaphase plate, with one chromosome facing each pole
- Microtubules from one pole are attached to the kinetochore of one chromosome of each tetrad
- Microtubules from the other pole are attached to the kinetochore of the other chromosome

Anaphase I

- In anaphase I, pairs of homologous chromosomes separate
- One chromosome moves toward each pole, guided by the spindle apparatus
- Sister chromatids remain attached at the centromere and move as one unit toward the pole

Telophase I and Cytokinesis

- In the beginning of telophase I, each half of the cell has a haploid set of chromosomes; each chromosome still consists of two sister chromatids
- Cytokinesis usually occurs simultaneously, forming two haploid daughter cells


بنفس الوقت

- In animal cells, a cleavage furrow forms; in plant cells, a cell plate forms
- No chromosome replication occurs between the end of meiosis I and the beginning of meiosis II because the chromosomes are already replicated

- Division in meiosis II also occurs in four phases
 - ① Prophase II
 - ② Metaphase II
 - ③ Anaphase II
 - ④ Telophase II and cytokinesis
- Meiosis II is very similar to mitosis

Figure 13.8b

- OCCUR in sex cells/sex organs .



تصطف ال chromosomes
على ال metaphase plate
من ال homologous

During another round of cell division, the sister chromatids finally separate; four haploid daughter cells result, containing unduplicated chromosomes.

Haploid daughter cells forming

genetic variation

Prophase II

- In prophase II, a spindle apparatus forms
- In late prophase II, chromosomes (each still composed of two chromatids) move toward the metaphase plate

Metaphase II

- In metaphase II, the **sister chromatids** are arranged at the metaphase plate
- Because of crossing over in meiosis I, the two sister chromatids of each chromosome are no longer genetically identical
- The kinetochores of sister chromatids attach to microtubules extending from opposite poles

Anaphase II

- In anaphase II, the sister chromatids separate
- The sister chromatids of each chromosome now move as two newly individual chromosomes toward opposite poles

Telophase II and Cytokinesis


- In telophase II, the chromosomes arrive at opposite poles
- Nuclei form, and the chromosomes begin decondensing

- Cytokinesis separates the cytoplasm
- At the end of meiosis, there are four daughter cells, each with a haploid set of unreplicated chromosomes
- Each daughter cell is genetically distinct from the others and from the parent cell

A Comparison of Mitosis and Meiosis

- ^{يحافظ} Mitosis conserves the number of chromosome sets, producing cells that are genetically identical to the parent cell
- ^{يقلل} Meiosis reduces the number of chromosomes sets from two (diploid) to one (haploid), producing cells that differ genetically from each other and from the parent cell

Figure 13.9



SUMMARY		
Property	Mitosis	Meiosis
DNA replication	Occurs during interphase <u>before mitosis begins</u>	Occurs during interphase <u>before meiosis I begins</u>
Number of divisions	One, including prophase, metaphase, anaphase, and telophase	Two, each including prophase, metaphase, anaphase, and telophase
Synapsis of homologous chromosomes	Does not occur	Occurs during prophase I along with crossing over between nonsister chromatids; resulting chiasmata hold pairs together due to sister chromatid cohesion
Number of daughter cells and genetic composition	Two, each diploid ($2n$) and genetically identical to the parent cell	Four, each haploid (n), containing half as many chromosomes as the parent cell; genetically different from the parent cell and from each other
Role in the animal body	Enables multicellular adult to arise from zygote; produces cells for growth, repair, and, in some species, asexual reproduction	Produces gametes; reduces number of chromosomes by half and introduces genetic variability among the gametes

① development

②

③

④ sexual reproduction

Figure 13.9a




Figure 13.9b

SUMMARY		
Property	Mitosis	Meiosis
DNA replication	Occurs during interphase before mitosis begins	Occurs during interphase before meiosis I begins
Number of divisions	One, including prophase, metaphase, anaphase, and telophase	Two, each including prophase, metaphase, anaphase, and telophase
Synapsis of homologous chromosomes	Does not occur	Occurs during prophase I along with crossing over between nonsister chromatids; resulting chiasmata hold pairs together due to sister chromatid cohesion
Number of daughter cells and genetic composition	Two, each diploid ($2n$) and genetically identical to the parent cell	Four, each haploid (n), containing half as many chromosomes as the parent cell; genetically different from the parent cell and from each other
Role in the animal body	Enables multicellular adult to arise from zygote; produces cells for growth, repair, and, in some species, asexual reproduction	Produces gametes; reduces number of chromosomes by half and introduces genetic variability among the gametes

- Three events are unique to meiosis, and all three occur in meiosis I
 - ① Synapsis and crossing over in prophase I: Homologous chromosomes physically connect and exchange genetic information
 - ② At the metaphase plate, there are paired homologous chromosomes (tetrads), instead of individual replicated chromosomes
 - ③ At anaphase I, it is homologous chromosomes, instead of sister chromatids, that separate

- Sister chromatid cohesion allows sister chromatids of a single chromosome to stay together through meiosis I
- Protein complexes called cohesins are responsible for this cohesion
- In mitosis, cohesins are cleaved at the end of metaphase
- In meiosis, cohesins are cleaved along the chromosome arms in anaphase I (separation of homologs) and at the centromeres in anaphase II (separation of sister chromatids)

Concept 13.4: Genetic variation produced in sexual life cycles contributes to evolution

- Mutations (changes in an organism's DNA) are the original source of genetic diversity
- Mutations create different versions of genes called alleles
- ^{إعادة تشكيل} Reshuffling of alleles during sexual reproduction produces genetic variation

Origins of Genetic Variation Among Offspring

- The behavior of chromosomes during meiosis and fertilization is responsible for most of the variation that arises in each generation
- Three mechanisms contribute to genetic variation
 - ① Independent assortment of chromosomes
 - ② Crossing over
 - ③ Random fertilization



مستقل


توزیع

Independent Assortment of Chromosomes

- Homologous pairs of chromosomes orient randomly at metaphase I of meiosis
- In independent assortment, each pair of chromosomes sorts maternal and paternal homologs into daughter cells independently of the other pairs

- The number of combinations possible when chromosomes assort independently into gametes is 2^n , where n is the haploid number
- For humans ($n = 23$), there are more than 8 million (2^{23}) possible combinations of chromosomes

Possibility 1



**Two equally probable
arrangements of
chromosomes at
metaphase I**

Possibility 2




Figure 13.10-2




Figure 13.10-3



*2 paternal
2 maternal*

2^n

$2^2 = 4$

Possibility 1



Possibility 2



Two equally probable arrangements of chromosomes at **metaphase I**

- The homologous pairs of chromosomes, each pair will align independently from each other (they are independently assorted)

Metaphase II



Daughter cells

Combination 1 **Combination 2**

Combination 3 **Combination 4**

$$2n = 4$$

$$n = 2$$

② *Crossing Over*

- Crossing over produces **recombinant chromosomes**, which combine DNA inherited from each parent
- Crossing over begins very early in **prophase I**, as homologous chromosomes pair up gene by gene


أجزاء

- In crossing over, homologous portions of two nonsister chromatids trade places
- Crossing over contributes to genetic variation by combining DNA from two parents into a single chromosome

Figure 13.11-1

Prophase I of meiosis

Pair of homologs



Nonsister chromatids
held together
during synapsis

Figure 13.11-2

Prophase I of meiosis




Figure 13.11-3

Prophase I of meiosis





Figure 13.11-4

Prophase I of meiosis


Pair of homologs

Nonsister chromatids held together during synapsis



TEM

Anaphase I



Anaphase II



Figure 13.11-5





Figure 13.11a



© 2011 Pearson Education, Inc.

3 *Random Fertilization*

- Random fertilization adds to genetic variation because any sperm can fuse with any ovum (unfertilized egg) نو بیج!
- The fusion of two gametes (each with 8.4 million possible chromosome combinations from independent assortment) produces a zygote with any of about 70 trillion diploid combinations

- Crossing over adds even more variation
- Each zygote has a unique genetic identity

• أهمية الـ genetic variation : تنتج صفات جديدة .





Animation: Genetic Variation

The Evolutionary Significance of Genetic Variation Within Populations

- Natural selection results in the accumulation of genetic variations favored by the environment
- Sexual reproduction contributes to the genetic variation in a population, which originates from mutations

Figure 13.12





Prophase I: Each homologous pair undergoes synapsis and crossing over between nonsister chromatids with the subsequent appearance of chiasmata.

Metaphase I: Chromosomes line up as homologous pairs on the metaphase plate.

Anaphase I: Homologs separate from each other; sister chromatids remain joined at the centromere.

Figure 13.UN02




Figure 13.UN03





Figure 13.UN04



The chromosomes of one color make up a haploid set.
All red and blue chromosomes together make up a diploid set.