



Pulmonary Gas Diffusion

Physical principles of gas exchange

Dr. Waleed R. Ezzat

وَقُلْ رَبِّ زِدْنِي عِلْمًا



RESPIRATORY SYSTEM

HA4AT BATCH



SUBJECT : Physiology

LEC NO. : 6

DONE BY : Sanaad Haitham

Lecture Objectives:

1. Review Dalton's law & Henry's law and understand their application to partial pressure of gasses in airways and blood.
2. Identify the values of PO_2 and PCO_2 in inspired air, alveolar air, and expired air. And identify the PO_2 and PCO_2 of arterial and venous blood.
3. Discuss and describe the diffusion of O_2 and CO_2 through the alveolar capillary membrane and list the factors affecting the rate of gas diffusion (Fick's law of diffusion).
4. Distinguish between perfusion limitation and diffusion limitation of gas transfer in the lung.
5. Define oxygen diffusing capacity and describe the rationale and technique for the use of carbon monoxide to determine diffusing capacity.
6. List main causes leading to decreased diffusion capacity of the lung.

Physics of Gas Diffusion and Gas Partial Pressures:

1. The Simple diffusion of a gas occurs from the high-concentration area toward the low-concentration area (i.e. it follows the concentration gradient). In a mixture of gases the rate of diffusion of each gas in the mixture is directly proportional to the **partial pressure** of that gas.

لها للتذكير الضغط الجزئي للغاز
= نسبة الغاز \times الضغط الكلي للغازات

2. The partial pressures of individual gases in a mixture are designated by the symbols PO_2 , PCO_2 , PN_2 , PHe , and so forth. The same symbols are used to describe the partial pressure of dissolved gases.

$P_{aO_2} = P_{AO_2}$ ✖
لأن الدم الخارج من الشعيرة
يذهب مباشرة من الـ Capillary إلى
Pulmonary Veins ثم Right atrium
✖ P : Pressure P_a : arterial Pressure
artery ← right ventricle
 P_A : Alveolar Pressure

Physics of Gas Diffusion and Gas Partial Pressures (cont.):

* نلاحظ أن الأوكسجين لا يمتص
عدد كبير من الجزيئات الذاتية لزيادة الضغط الجزئي

3. The number of dissolved molecules of a gas depends on the **solubility coefficient** of that gas. A gas with high water solubility, like CO_2 , large number of molecules can be dissolved without building up excess partial pressure within the solution. Conversely, in the case of molecules with low solubility, like O_2 , high partial pressure will develop with fewer dissolved molecules (**Henry's law**).

$$\text{Partial pressure} = \frac{\text{Concentration of dissolved gas}}{\text{Solubility coefficient}}$$

لافتة كل ما زادته قابلية الغاز للذوبان قلت الضغط

*
فصل

The concentration is expressed in volume of gas dissolved in each volume of water.

The solubility coefficients for important respiratory gases at body temperature

Oxygen	0.024	
Carbon dioxide	0.57	
Carbon monoxide	0.018	1. قريب من غاز الأوكسجين لذلك يمكن إجراء الأبحاث والتجارب عليه في الأن بعد ما يقبر من
Nitrogen	0.012	الهويصلات إلى الدم ينتقل كل إلى الـ (RBCs) ولا يذهب جزء منه للبلازما فتكون الحسابات أبسط في
Helium	0.008	2. نأخذ بعينه الاعتبار فارق الأرقام وسمية الغاز

لهذا السبب يمكن نقل O_2 زائداً في البلازما أما CO_2 لا يمكن نقله إلا بالـ (HB).

Note: CO_2 is more than 20 times as soluble as O_2

Physics of Gas Diffusion and Gas Partial Pressures (cont.):

	Alveoli	Venous blood	
P_{O_2}	100	40	الأكسجين
P_{CO_2}	40	46	ثاني أكسيد الكربون
الضغط	mm hg		

4. In the alveolus, the direction of diffusion of a gas is determined by the difference of the partial pressures of that gas across the alveolar membrane. For the O_2 , it diffuses and dissolves into the blood as its partial pressure is greater in the alveolar air. Alternatively, the net diffusion of CO_2 occurs toward the alveolar air as its partial pressure is greater in the dissolved state in the blood.

طبقاً
لأن كمية
ال O_2
رح تزيد !!

تكون أكثر
في الدم
exchange blood في

5. At normal body temperature, $37^\circ C$, the partial pressure of water vapor in the alveolar gas mixture is **47 mmHg**. It is designated as PH_2O .



ضغط بخار الماء

Physical factors affecting the rate of gas diffusion (Fick's law)

... \propto طردي direct $\uparrow \uparrow$
... \propto عكسي inverse $\uparrow \downarrow$

1. The partial pressure difference (direct relationship)
2. The solubility of the gas in the fluid (direct relationship). The greater the solubility of the gas, the greater the number of molecules available to diffuse for any given partial pressure difference.
3. The cross-sectional area of the fluid or surface area of the respiratory membrane (direct relationship). The greater the cross-sectional or surface area of the diffusion pathway, the greater the total number of molecules that diffuse decreases. Emphysema and removal of an entire lung decreases the total surface area of the respiratory membrane.
4. The distance through which the gas must diffuse (inverse relationship). The greater the distance the molecules must diffuse, the longer it will take the molecules to diffuse the entire distance. Some pulmonary diseases cause fibrosis of the lungs and can increase the thickness of some portions of the respiratory membrane (\uparrow distance) \rightarrow impair gaseous exchange.

الأمراض التي تزيد سمك الحاجز التنفسي بين
الأكياس (alveoli) والدم (capillaries) \rightarrow impair gaseous exchange.

Physical factors affecting the rate of gas diffusion (cont.)

5. *المسألة في الوزن الجزيئي* The molecular weight of the gas (inverse relationship). The kinetic movement of the molecules is inversely proportional to the square root of the molecular weight. The greater the velocity of kinetic movement, the greater the rate of diffusion of the gas.
6. *تعدل لانتفا ثابتة* The temperature of the fluid (direct relationship). This factor does not need to be considered since body temperature remains reasonably constant.

في حالة الجسم تتساوى لجميع الغازات باستثناء (MW و S)

$$\text{Diffusion rate (D)} \propto \frac{\Delta P \times A \times S}{d \times \sqrt{MW}}$$

ΔP is the partial pressure difference between the two ends of the diffusion pathway, **A** is the cross-sectional area of the pathway, **S** is the solubility of the gas, **d** is the distance of diffusion, and **MW** is the molecular weight of the gas.

Physical factors affecting the rate of gas diffusion (cont.)

Note: سرعة انتشار الغاز
من طرية (alveolar membrane)

1. The **diffusion coefficient** of the gas is determined by **two** factors, namely, S/\sqrt{MW} . This means that the relative rates at which different gases at the same partial pressure levels will diffuse are proportional to their diffusion coefficients.
تغير ΔP
2. If the diffusion coefficient for O_2 is assumed as 1, then, the relative diffusion coefficient for CO_2 is 20.3. Therefore, for a given pressure difference, CO_2 diffuses about 20 times as rapidly as O_2 .
* إذا كان هناك مشكلة في (alveolar membranes) يتأثر مستوى الـ (O_2) بشكل أكبر لأن
3. Since both oxygen and carbon dioxide are **highly lipid soluble**, then the major limitation to the movement of these gases in tissues is the rate at which the gases can diffuse through the tissue water instead of through the cell membranes.
لا ببساطة هذه الغازات (lipid-soluble) / تعيد الماء داخل الـ (tissue) بمعدية
تغير الـ (Cell membrane) بسهولة

مهم جدًا في حفظ الحد.

Partial Pressures of Respiratory Gases (in mm Hg) as They Enter and Leave the Lungs (at Sea Level)

	Atmospheric Air	Humidified Air	Alveolar Air	Expired Air
N ₂	597 (78.62)	563.4 (74.09)	569 (74.9)	566 (74.5)
O ₂	159 (20.84)	149.3 (19.67)	100 ← 104 (13.6)	120 (15.7)
CO ₂	0.3 (0.04) ← 0	0.3 (0.04)	40 (5.3)	27 (3.6)
H ₂ O	3.7 (0.50)	47 (6.20)	47 (6.2)	47 (6.2)
Total	760 (100)	760 (100)	760 (100)	760 (100)

$$\begin{array}{l}
 160 \rightarrow 40 \\
 100 = 100 \\
 46 \rightarrow 0 \\
 40 = 40
 \end{array}$$

* داخل الجسم (P_{O_2}) قبل التنفس 40 mmHg

والهواء الخارجي 160 mmHg عند وصوله إلى

الحويصلة يتساوى (P_{O_2}) الحويصلة والدم = 100 mmHg

✗ داخل الجسم (P_{CO_2}) قبل التنفس 46 mmHg

والهواء الخارجي قريب الصفر لكنه ينتقل بكفاءة عالية للخارج

فيصبح 40 mmHg داخل الدم وفي الحويصلة تحل هذه النسبة كل فروع الهواء تدريجيًا لأن P_{CO_2} قليل في الهواء الخارجي.

Oxygen concentration and P_{AO_2}

التذكير يمكن زيادة الـ (alveolar ventilation) إلى (45 l/min)

زاد أربعة أضعاف (O₂ absorption) يجب أن تزيد الـ (ventilation) أربع أضعاف للحفاظ على (P_{O₂})

- P_{AO_2} is determined by (1) the rate of absorption of O₂ into the blood and (2) the rate of entry of new O₂ into the lungs by the ventilatory process.


- **Red curve** → O₂ absorption at a rate of 250 ml/min (rest level)

- **Blue curve** → O₂ absorption at a rate of 1000 ml/min (exercise)

- Note that the rate of alveolar ventilation increases fourfold to maintain the alveolar PO₂ at the normal value of **104 mmHg**.


رقم مقدس
نحافظ عليه

أضعاف



Carbone dioxide concentration and P_{ACO_2}

- P_{ACO_2} is determined by (1) the rate of excretion of CO_2 into the blood and (2) the rate of removal of CO_2 from the alveoli by ventilation.
- **Red curve** → normal rate of CO_2 excretion of 200 ml/min (rest level)
- **Blue curve** → CO_2 excretion rate of 800 ml/min (exercise)
- Note that (1) the alveolar PCO_2 increases directly in proportion to the rate of CO_2 excretion, as represented by the fourfold elevation of the curve, and (2) the alveolar PCO_2 decreases in inverse proportion to alveolar ventilation. The operating point for alveolar PCO_2 is at point A in the figure (i.e., 40 mmHg).



نفس السلاير السابق

لكن مكي (P_{CO_2})

الأرقام ليست دقيقة
المفروض (20 L/min)

رقع مقدس

Diffusing Capacity of the Respiratory Membrane

- The diffusion capacity of a gas is the volume of that gas that will diffuse through the membrane each minute for a partial pressure difference of 1 mmHg.

كمية الغاز التي تتحرك
عندما يكون (الغاز P) = 1 mmHg

- The time courses for the red blood cell to move through the pulmonary capillary takes about 0.75 s.


وقت قدوم
الدم من الشعيرة

- Under typical resting conditions, the capillary PO_2 virtually reaches that of alveolar gas when the red cell is about one-third of the way along the capillary. Therefore, O_2 transfer (from alveolus to blood) is **perfusion limited** and not diffusion limited.

تصلح
ثلث الوقت
فقط لتبادل
الغازات.

- The **diffusing capacity for O_2** under resting conditions averages **21 ml/min/mmHg**. This capacity increases during exercise.

ال (Perfusion) قليلة أو ال (Perfusion) قليل (يقل ال O_2 transfer).



ال Diffusion سريع جداً
لن يتأثر مهما كانت كمية

ال (Perfusion) قليلة أو ال (Perfusion) قليل (يقل ال O_2 transfer).

Diffusing Capacity of the Respiratory Membrane (cont.)

التجربة التي طالعها فيها
الأرقام

- With severe exercise, the pulmonary blood flow is greatly increased, and the time spent by the red cell in the capillary may be reduced to as little as 0.25 s. However, in normal subjects breathing air, there is still no measurable fall in end-capillary PO_2 .

Since the diffusion coefficient of CO_2 is slightly more than 20 times that of O_2 , it can be assumed that a diffusing capacity for CO_2 under resting conditions is about 400 to 450 ml/min/mmHg and during exercise of about 1200 to 1300 ml/min/mmHg.

Physiologists usually measure carbon monoxide (CO) diffusing capacity instead of O_2 diffusing capacity as it is easier. Oxygen diffusing capacity is obtained by multiplying CO diffusing capacity by a factor of 1.23. The average diffusing capacity for CO in healthy young men at rest is 17 ml/min/mmHg.




Figure Diffusing capacities for carbon monoxide, oxygen, and carbon dioxide in the normal lungs under resting conditions and during exercise.

التجربة استخدم
فيها (CO) فليس معامل
تصحيح لتصحيح الأرقام.

يُسْتَقَرَّت
ثُلُثُ
مُتَغَيَّرَةِ
الْزَّمَنِ
تُغَيَّرُ
الْشِدَّةِ

بسبب
الزيادة
يُنْجِزُ بِسُرْعَةٍ
فِي الْمَاءِ
عَلَى حِدَارٍ

الأنظمة
تُغَيَّرُ
بِسُرْعَةٍ

Test Question:

Q. Concerning the diffusing capacity of the lung:

- A. It is best measured with carbon monoxide because this gas diffuses very slowly across the blood-gas barrier.
- B. Diffusion limitation of oxygen transfer during exercise is more likely to occur at sea level than at high altitude.
- C. Breathing oxygen reduces the measured diffusing capacity for carbon monoxide compared with air breathing.
- D. It is decreased by exercise.
- E. It is increased in pulmonary fibrosis, which thickens the blood-gas barrier.

Answer : C

يصبح الارتباط (C) بالهيموغلوبين
صعباً لأنه يتنافس (O₂) فقط جزئياً ترتبط
بالهيموغلوبين (C₂)