

imp

Contraction of smooth muscles


Begins when intracellular Ca^{2+} increases

① بداية ال Contraction في ال smooth muscle كذا زيادة ال intracellular Ca^{2+} لكن بظروف مختلفة عن ال skeletal muscle

Intracellular calcium ion (Ca^{++}) concentration increases when Ca^{++} enters the cell through calcium channels in the cell membrane or is released from the sarcoplasmic reticulum.

The Ca^{++} binds to calmodulin (CaM) to form a Ca^{++} - CaM complex, which then activates myosin light chain kinase ($MLCK$).

The active $MLCK$ phosphorylates the myosin light chain leading to attachment of the myosin head with the actin filament and contraction of the smooth muscle.



↓ Ca^{+2} → 3 ways to ↓ intracellular Ca^{+2} (check pic)


↓
deactivation →
MLCK
↓
dephosphorylation

Imp

Relaxation of smooth muscle


Relaxation occurs when calcium ion (Ca^{++}) concentration decreases below a critical level Ca^{++} is pumped out of the cell or into the sarcoplasmic reticulum.

Ca^{++} is then released from calmodulin (CaM) and myosin phosphatase removes phosphate from the myosin light chain, causing detachment of the myosin head from the actin filament and relaxation of the smooth muscle.



Sequence of events in contraction and relaxation of smooth muscle.

Acetylcholine is given as a source of stimulus in this example



Properties of smooth muscle contraction

- Low Energy Requirement to sustain smooth muscle Contraction and low O₂ consumption
- Can operate over large range of lengths (60 - 75% shortening possible)
- length tension relation ship is over a wide range
- Can be **myogenic** (spontaneously active) (pre-muscle)
- Has **Ca²⁺ action potentials**. Ca entering through channels is a very important source of calcium for contraction
- Smooth muscles exhibits sustained prolonged tonic contraction which may last for hours or even days
- **Grading of muscle contraction**
 - Depends on intracellular Ca Ions concentration ,
 - No recruitment, specially in visceral smooth muscles

از Ca²⁺ لور
المسؤول عن
الانقباض (depolarization)

Contraction
سعة طويلة

ال tension يعتمد على
Ca²⁺ + length

تدرج في زيادة
اد tension

Summation في Ca²⁺

Characteristics of smooth muscle contraction

4

1 Slow muscle contraction

- Slow contraction due to slow cycling of cross bridges
- Degradation of the ATP that energizes the movements of the cross-bridge heads is greatly reduced, with corresponding slowing of the rate of cycling.
- longer latency to respond to stimulus
- Latency : begins to contract 50 to 100 milliseconds after it is excited
- initiation of contraction in response to calcium ions is much slower than in skeletal muscle
- Reaches full contraction about 0.5 second later, and then declines in contractile force in another 1 to 2 seconds
- Total contraction time of 1 to 3 seconds.
- This is about 30 times as long as a single contraction of an average skeletal muscle fiber.
- Other types of muscle contraction could be as short as (0.2 sec - 30 sec)

• Ca^{+2} storage is supplemented by caveolae [small vesicles that cluster close to cell membrane]

Properties of smooth muscle contraction

Long duration of contraction



• Slow Cycling of the Myosin Cross-Bridge

⇒ Because of the slow ATPase activity

- attachment to actin, then release from the actin, and reattachment for the next cycle—is much slower than in skeletal muscle
- The fraction of time that the cross-bridges remain attached to the actin filaments, which is a major factor that determines the force of contraction, is believed to be greatly increased in smooth muscle.
- A possible reason for the slow cycling is that the cross-bridge heads have far less ATPase activity than in skeletal muscle; thus, degradation of the ATP that energizes the movements of the cross-bridge heads is greatly reduced, with corresponding slowing of the rate of cycling

③

• **Low Energy Requirement to Sustain Smooth Muscle Contraction** (energy efficient)

دیت O_2 كى كۆپىنچە

- Only 1/10 to 1/300 as much energy is required to sustain the same tension of contraction in smooth muscle as in skeletal muscle. This, too, is believed to result from the slow attachment and detachment cycling of the cross-bridges, and because only one molecule of ATP is required for each cycle, regardless of its duration
- This low energy utilization by smooth muscle is important to the overall energy economy of the body because organs such as the intestines, urinary bladder, gallbladder, and other viscera organs

4

• **Slowness of Onset of Contraction and Relaxation of the Total Smooth Muscle Tissue**

- Longer latency than that of skeletal muscle after it is excited, reaches full contraction about 0.5 second later, and then declines in contractile force in another 1 to 2 seconds, giving a total contraction time of 1 to 3 seconds.
- This is about 30 times as long as a single contraction of an average skeletal muscle fiber. However, because there are so many types of smooth muscle, contraction of some types can be as short as 0.2 second or as long as 30 seconds.
- The slow onset of contraction of smooth muscle, as well as its prolonged contraction, is caused by the slowness of attachment and detachment of the cross-bridges with the actin filaments. In addition, the initiation of contraction in response to calcium ions is much slower than in skeletal muscle,

Comparison of muscle twitch in different types of muscle


AP → contraction → relaxation

Duration of contraction

Skeletal < Cardiac < smooth

Speed of contraction

Smooth > Cardiac > skeletal



Smooth > skeletal

- **Maximum Force of Contraction Is Often Greater in Smooth Muscle Than in Skeletal Muscle.**
- Despite the relatively few myosin filaments in smooth muscle, and despite the slow cycling time of the cross-bridges, the maximum force of contraction of smooth muscle is often greater than that of skeletal muscle, as much as 4 to 6 kg/ cm² cross-sectional area for smooth muscle in comparison with 3 to 4 kilograms for skeletal muscle.
- This great force of smooth muscle contraction results from the prolonged period of attachment of the myosin cross-bridges to the actin filaments.

Numbers are insignificant!

سفنكتر العضلات
sphincter muscles

Latch Mechanism Facilitates Prolonged Holding of Contractions of Smooth Muscle

سبب توتره ال
مستمر في
عضلات
مخاطية
العضلة
مخاطية

• Smooth muscle can maintain tension for long periods
(Tonic contraction)

long time

• The latch state seems to occur because the cross bridges **do not** dissociate very rapidly in spite of the fact that the myosin light chain is dephosphorylated ;

• Accordingly energy expenditure is minimal

• This is thought to be important in sphincter muscles ex in Stomach where tension development must occur for long periods of time.

سؤال

Stress-Relaxation and reverse Relaxation in Smooth Muscle

- Mainly occurs in visceral ^{single unit} unitary type of smooth muscle of many hollow organs *specially the urinary bladder.*
- **Stress relaxation** \longrightarrow *عشان نقلل ال tension* *قابلية التمدد* *عندما يكون في ضغط (stress)*
- is its ability to return to nearly its original *force* of contraction seconds or minutes after it has been elongated or shortened.
- Response to stretch briefly, then return to their normal state of tension and adapt to new length
- For example, a sudden increase in fluid volume in the urinary bladder, thus stretching the smooth muscle in the bladder wall, causes an immediate large increase in pressure in the bladder.
- However, during the next 15 seconds to a minute or so, despite continued stretch of the bladder wall, the pressure returns almost exactly back to the original level Then, when the volume is increased by another step, the same effect occurs again.

Stress-Relaxation and reverse Relaxation in Smooth Muscle

- Reverse stress-relaxation *micturition*
- When the volume is suddenly decreased, the pressure falls drastically at first but then rises in another few seconds or minutes to or near the original level.
- These phenomena are called stress-relaxation and and reverse stretch relaxation , they allow a hollow organ to maintain about the same amount of pressure inside its lumen despite sustained large changes in volume.

إرخاع
Stress relaxation

الامتداد: مع زيادة الحجم يبقى الضغط (tension) ثابتة

Effects of Local Tissue Factors on smooth muscle contraction

- Lack of ^{hypoxia} oxygen in the local tissues causes smooth muscle relaxation and, therefore, vasodilation.
- Excess carbon dioxide causes vasodilation. 3. Increased hydrogen ion concentration causes vasodilation.
- Adenosine, lactic acid, increased potassium ions, nitric oxide, and increased body temperature can all cause local vasodilation.
- Decreased blood pressure, by causing decreased stretch of the vascular smooth muscle, also causes these small blood vessels to dilate.

نقصان ال pH

Effects of Hormones on Smooth Muscle Contraction

- Many circulating hormones in the blood affect smooth muscle contraction to some degree, and some have profound effects.
- Among the more important of these hormones are norepinephrine, epinephrine, angiotensin II, endothelin, vasopressin, oxytocin, serotonin, and histamine.
- A hormone causes contraction of a smooth muscle when the muscle cell membrane contains hormone-gated excitatory receptors for the respective hormone. Conversely, the hormone causes inhibition if the membrane contains inhibitory receptors for the hormone rather than excitatory receptors.

Mechanisms of Smooth Muscle Excitation or Inhibition by Hormones or Local Tissue Factor

- Excitation
- Some hormone receptors in the smooth muscle membrane open sodium or calcium ion channels and depolarize the membrane, the same as after nerve stimulation.
- Depolarization allows for calcium ion entry into the cell, which promotes the contraction.
- Inhibition
- Occurs when the hormone (or other tissue factor) closes the sodium and calcium channels to prevent entry of these positive ions; inhibition also occurs if the normally closed potassium channels are opened, allowing positive potassium ions to diffuse out of the cell. Both these actions increase the degree of negativity inside the muscle cell, a state called hyperpolarization, which strongly inhibits muscle contraction

Other mechanism of hormones affecting smooth muscle contraction : Role of cAMP and cGMP

- To inhibit contraction, other receptor mechanisms are known to activate the enzyme adenylate cyclase or guanylate cyclase in the cell membrane. The portions of the receptors that protrude to the interior of the cells are coupled to these enzymes, causing the formation of cyclic adenosine monophosphate (cAMP) or cyclic guanosine monophosphate (cGMP), so-called second messengers.
- cAMP or cGMP has many effects, one of which is to change the degree of phosphorylation of several enzymes that indirectly inhibit contraction. The pump that moves calcium ions from the sarcoplasm into the sarcoplasmic reticulum is activated, as well as the cell membrane pump that moves calcium ions out of the cell; these effects reduce the calcium ion concentration in the sarcoplasm, thereby inhibiting contraction.

MEMBRANE POTENTIALS AND ACTION POTENTIALS IN VISCERAL SMOOTH MUSCLE

single unit SM


- In the normal resting state, the intracellular potential is usually about -50 to -60 millivolts \Rightarrow Resting membrane potential for single-unit SM
- The action potentials of visceral smooth muscle occur in one of two forms

- ① spike potentials
- ② action potentials with plateau

- Such action potentials can be elicited in many ways

- Electrical stimulation.
- Spontaneous generation in the muscle fiber \rightarrow pacemaker activity
- Hormones
- transmitter substances from nerve fibers
- Stretch

5 factors can elicit AP in single-unit smooth muscles




Spike potentials \leftarrow يشوفونها لوحدها أو مع slow waves

- Typical smooth muscle action potential (spike potential) elicited by an external stimulus.
- Observed in GT and in most types of unitary smooth muscle.
- The duration of this type of action potential is 10 to 50 milliseconds, *زمنه الـ skeletal m*
- Such action potentials can be elicited in many ways—for example, by electrical stimulation, by the action of hormones on the smooth muscle, by the action of transmitter substances from nerve fibers, by stretch, or slow waves depolarization \rightarrow NE or ACh
- Sodium participates little in the generation of the action potential in most smooth muscle. \rightarrow دور الـ Na في الـ smooth muscle هو تقال الـ AP الى الـ threshold
- Instead, flow of calcium ions to the interior of the fiber is mainly responsible for the action potential *طبيعته بيتسبب من الـ Ca²⁺ الـ Depolarization !!*
- Depolarization phase is mainly due to activation of L type Ca channel *Ca²⁺ influx*
- Receptive spike potentials generated by slow wave depolarization

كيف ممكن يهبط حنا AP بسبب الـ stretch??

الـ smooth muscle الـ stretch يفتح الـ channels الـ Na⁺ influx \rightarrow Depolarization \rightarrow threshold \rightarrow AP



يكون في الخداج وفي غيره الـ pacemaker activity

Spike potential with slow waves

Spike potential alone

Spike AP

Slow waves

Re (K⁺ efflux)

Ca²⁺ influx

Ca²⁺ influx

Pumpy channels:

قنوات دالها مفتوحة الـ Ca²⁺ و الـ Na⁺

و الـ membrane بغير أكثر




Action potential with a plateau, recorded from a smooth muscle fiber of the uterus

The onset of this action potential is similar to that of the typical spike potential.

However, instead of rapid repolarization of the muscle fiber membrane, the repolarization is delayed for several hundred to as much as 1000 milliseconds (1 second).


The importance of the plateau is that it can account for the prolonged contraction that occurs in some types of smooth muscle, such as the ureter, the uterus under some conditions, and certain types of vascular smooth muscle.



Slow Wave Potentials (*pacemaker waves*) in Unitary Smooth Muscle Can

- action potentials arise within the smooth muscle cells **without an extrinsic stimulus** (بدون محفز خارجي)
- action potentials arise within the smooth muscle cells without an extrinsic stimulus
- Slow wave depolarization Leads to Spontaneous Generation of Action Potentials. (spike potentials)
- **the slow waves are caused by waxing and waning of the pumping of Na pump.** Another suggestion is that the conductances of the ion channels increase and decrease rhythmically
- This type of pacemaker activity for example in the gut
- Controls the rhythmical contractions of the gut.
- Slope of depolarization is influenced by ANS


of slow waves potentials



Single unit

Slow wave and spikes potentials in visceral smooth muscles Effects of sympathetic and parasympathetic stimulation

تثبيط ← why? ← تحفيز
لماذا يكون لدينا بنكري عن الأعضاء الداخلية أي يتم تثبيط عملها في الـ SNS وتحفيز عملها في الـ PSNS



Stimulation by
1. Norepinephrine
2. Sympathetics

Stimulation by
1. Stretch
2. Acetylcholine
3. Parasympathetics

Stimulation does NE الـ Smooth muscle receptors α receptors

bb slope
Inhibition (relaxation) or decreased contraction

العلاوة عوامل أي بتحفيز الـ slow waves